

University of Groningen

Picosecond Fluorescence of Intact and Dissolved PSI-LHCI Crystals

van Oort, Bart; Amunts, Alexey; Borst, Jan Willem; van Hoek, Arie; Nelson, Nathan; van Amerongen, Herbert; Croce, Roberta

Published in:
Biophysical Journal

DOI:
[10.1529/biophysj.108.140467](https://doi.org/10.1529/biophysj.108.140467)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van Oort, B., Amunts, A., Borst, J. W., van Hoek, A., Nelson, N., van Amerongen, H., & Croce, R. (2008). Picosecond Fluorescence of Intact and Dissolved PSI-LHCI Crystals. *Biophysical Journal*, 95(12), 5851-5861. <https://doi.org/10.1529/biophysj.108.140467>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Supplementary Material

Supplementary Figure Legends

Figure 9. Decay Associated Fluorescence Spectrum ($\tau = 6$ ps) of pinacyanol iodide in methanol at 294 K. Excitation was at 475 nm, fluorescence was detected with the streak-camera setup.


Figure 10. Sums of fluorescence decay curves of a PSI-LHCI crystal measured at room temperature with FLIM. Data were corrected for background signal, and normalized in the peak. Excitation was at 860nm, at energies as indicated, detection was at 670-730nm.

Figure 11. Decay Associated Spectra (DAS) of the slowest components of PSI-LHCI in solution at 287K, excited at 410nm and at 475nm (dashed), scaled on the total area of the 61 ps and 143 ps DAS (same data as in Figure 3 of the main text, but multiplied by 100).


Figure 12. Decay Associated Spectra (DAS) of PSI-LHCI in solution at 287K, upon excitation of only core or only LHCI pigments (dashed). DAS were calculated from linear combinations of the DAS in Figure 3 of the main text, with 65 % core excitation at 410 nm, and 30 % at 475 nm. See text for more details. DAS are scaled to equal area (by multiplication of DAS of 61 ps, 143 ps and 0.37 ns, respectively, by 34.9, 11.5 and 1049 for core excitation, and by 21.9, 67.9 and 670 for LHCI excitation).

Supplementary Figures


Supplementary Figure 9.



Supplementary Figure 10.



Supplementary Figure 11.



Supplementary Figure 12.

